



11/18/2013
UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

1/4

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

11-18-13

Stephen Adams / Erik Janus
Monsanto Company
1300 I (Eye) Street, NW
Suite 450 East
Washington, DC 20005

Subject: Label Amendment
EPA Reg. No.: 524-549 / RD 1617 Add Tank-Mix Products Use for Preemergence
Application in Soybean Supplemental Label

Dear Mr. Adams & Mr. Janus:

The supplemental labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable.

Submit one copy of the final printed label for the record before you release the product for shipment. A stamped copy of the label is enclosed for your records. The supplemental label must be incorporated into the master label within 18 months from the date of this letter. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions. If you have any questions please call Erik Kraft at 703-308-9358 or email at Kraft.Erik@epa.gov.

Sincerely,

Kable "Bo" Davis
Product Manager 25
Herbicide Branch
Registration Division (7505P)

SUPPLEMENTAL LABELING

This supplemental label expires on December 31, 2016 and must not be used or distributed after this date.


READ THE ENTIRE PRODUCT LABEL FOR [INSERT BRAND NAME] BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

When using [INSERT BRAND NAME] as permitted according to this supplemental labeling, read and follow all applicable directions, restrictions, and precautions on the product label provided with the pesticide container and on this supplemental labeling. This supplemental labeling must be in the possession of the user at the time of pesticide application.

RD 1617 Herbicide

Alternate Brand Names:
Roundup PowerMAX® Herbicide

EPA Reg. No. 524-549
[INSERT BRAND NAME] is a trademark of Monsanto Company



Use on Soybean

Keep out of reach of children

CAUTION!

In case of an emergency involving this product, call collect, day or night, 314-694-4000.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This labeling must be in the possession of the user at the time of herbicide application.

AVOID CONTACT OF THIS HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS, FRUIT OF CROPS, AND ANY DESIRABLE PLANTS OR TREES, EXCEPT AS DIRECTED FOR ROUNDUP READY® CROPS, AS SEVERE PLANT INJURY OR DESTRUCTION COULD RESULT.

See the "PRODUCT INFORMATION," "MIXING" and "ANNUAL AND PERENNIAL CROPS" sections of the [INSERT BRAND NAME] product label for important use information.

TYPES OF APPLICATION: Chemical Fallow; Preplant Fallow Beds; Preplant; At-Planting; Preemergence; Hooded Sprayer in Row Middles; Shielded Sprayer in Row Middles; Wiper Applicator in Row Middles; Post-Harvest

For directions for use with Roundup Ready soybean and Roundup Ready 2 Yield soybean, see the "ROUNDUP READY CROPS" section of the [INSERT BRAND NAME] product label.

Preplant, At-Planting, Preemergence

USE INSTRUCTIONS: This product may be applied alone or in a tank mixture before, during or after planting soybean, but prior to crop emergence.

TANK MIXTURES: This product may be tank-mixed with 2,4-D, Banvel or Clarity and applied prior to planting only. This product may be tank-mixed with the following products and applied prior to crop emergence. Ensure that the product being used is labeled for application prior to planting or the emergence of soybean. Read and follow label directions of all products in the tank mixture. Apply these tank mixtures in 10 to 20 gallons of water per acre.

[Insert active ingredient(s) or brand name of product(s) containing the following active ingredients that, at the time of printing, are registered for use preplant, at-planting and/or preemergence to soybean:

alachlor; atrazine; carfentrazone-ethyl; chlorimuron ethyl; clethodim; clomazone; cloransulam-methyl; dimethenamid; dimethenamid-p; fenoxypop; fluazifop-p-butyl; flufenacet; flumetsulam; flumiclorac pentyl ester; flumioxazin; fluthiacet-methyl; fomesafen; imazaquin; imazethapyr; lactofen; linuron; metolachlor; s-metolachlor; metribuzin; pendimethalin; pyroxasulfone; quizalofop P-ethyl; saflufenacil; sulfentrazone; tribenuron methyl; trifluralin

Aim; Assure II; Authority; Authority Assist; Authority First; Authority MAXX; Authority MTZ; Authority XL; Axiom; Blanket; Boundary; Cadet; Canopy; Canopy EX; Classic; Cobra; Command; Command Xtra; Dawn; Domain; Dual MAGNUM; Dual II MAGNUM; Fierce; FirstRate; Flexstar; Frontier; Fusion; Gangster; Gauntlet; INTRRO; Lexone; Linex; Linuron; Lorox; Lorox Plus; Me-Too-Lachlor; Micro-Tech; Optill; Outlook; PARRLAY; Pendimax; Phoenix; Prowl; Prowl H2O; Pursuit; Pursuit Plus; Python; Reflex; Resource; Rhythm; Scepter; Select; Select MAX; Sencor; Sharpen; Spartan; Squadron; Steel; Treflan; Valor; Valor XLT]

For tough-to-control annual weeds such as fall panicum, barnyardgrass, crabgrass, shattercane and broadleaf signalgrass up to 2 inches tall, and Pennsylvania smartweed up to 6 inches tall, apply 22 fluid ounces of this product per acre in a tank mixture with one of the products listed. For other annual weeds listed on the **[INSERT BRAND NAME]** product label, apply 16 to 22 fluid ounces of this product per acre when weeds are less than 6 inches tall and 22 to 32 fluid ounces per acre when weeds are over 6 inches tall.

Spot Treatment

USE INSTRUCTIONS: This product may be applied as a spot treatment prior to initial pod set in soybean.

RESTRICTIONS: Do not apply this product to more than 10 percent of the total field area to be harvested.

Selective Equipment

USE INSTRUCTIONS: This product may be applied in soybean using a shielded applicator, hooded sprayer, wiper applicator or sponge bar. See additional instructions on the use of selective equipment in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of the **[INSERT BRAND NAME]** product label.

RESTRICTIONS: Allow a minimum of 7 days between application and harvest.

Preharvest

USE INSTRUCTIONS: This product may be applied to soybean prior to harvest after pods have set and lost all green color. Apply at rates given in the "ANNUAL WEEDS RATE SECTION" and "PERENNIAL WEEDS RATE SECTION" of the **[INSERT BRAND NAME]** product label. Take care to avoid excessive seed shatter loss due to ground application equipment.

RESTRICTIONS: Do not apply more than 3.3 quarts of this product per acre for preharvest application using ground application equipment or more than 44 fluid ounces per acre when making the application using aerial application equipment. Allow a minimum of 7 days between application and harvest of soybean. If the preharvest application rate is greater than 22 fluid ounces per acre, do not graze or harvest hay or fodder within the application area for livestock feed within 25 days of application. If the application rate is 22 fluid ounces per acre or less, the grazing restriction is reduced to 14 days after application.

Read the "Limit of Warranty and Liability" in the product label booklet for **[INSERT BRAND NAME]** before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

© [YEAR]

4/4

MONSANTO COMPANY
800 N. LINDBERGH BLVD.
ST. LOUIS, MISSOURI 63167 USA

[INSERT DATE]

[PRINT PLATE NUMBER]